

# The Covid-19 *in vivo* Nanoparticle Vaccine

Thomas Prevenslik

QED Radiations, Berlin 10777 Germany

thomas@nanoqed.org

## 1. Abstract

Classical physics allows the atom to have heat capacity at the nanoscale, the conservation of heat proceeding by a change in temperature. However, simple QED based on the Planck law of quantum mechanics denies the atom in nanostructures the heat capacity to conserve heat by a change in temperature, the consequence of which is any heat is conserved by creating standing EM radiation that is released to the surroundings. Unlike electronic quantum states, simple QED is based on size dependent quantum states depending on the dimensions of the nanostructure over which the EM waves stand. UVC radiation is known to disinfect Coronavirus in the air or on surrounding surfaces, but not inside body organs. In this regard, patients diagnosed positive for Covid-19 are proposed treated by *in vivo* vaccinations of  $\sim 80$  nm lipid nanoparticles that emit UVC radiation powered only by body heat. In the blood stream, the NPs may enter the brain and damage neurons and DNA, but in brief NP treatments the risk of brain damage is expected to be minimal. CDC testing to determine acceptable NP doses is requested.

Keywords: Covid-19, Quantum Mechanics, Planck law, Nanoparticles.

## 2. Introduction

The disinfection of viruses by UV radiation [1] has a long history. In 1877, microorganisms in test tubes containing Pasteur's solution, an artificial nutrient fluid for cultivating organisms, upon exposure to sunlight prevented [2] the growth of the organisms for several months. About the same time, Tyndall confirmed [3] sunlight neutralized organisms dependent on intensity, duration, and wavelength, with the UV wavelengths of the solar spectrum being the most effective. However, the specific UV wavelengths of organism disinfection of about 260 nm were not identified [4] until 1944. Thereafter, the production of pyrimidine dimers in DNA that block virus reproduction was demonstrated after exposure to UV radiation, thereby providing the basis for UV disinfection of biological organisms.

In 2020, UV disinfection of viruses became of great importance. The Coronavirus known as Covid-19 became a pandemic that changed the economic future of the world, although in severity only comparable to influenza. Nevertheless, there is no vaccine or specific treatment known for Covid-19. In this regard, Coronavirus disinfection by external UV radiation sources would resolve the pandemic research, but only if the virus is still in the air or on the surface of surroundings.

In the usual Covid-19 infection, the virus has infected the blood stream and internal organs. External UV radiation no longer works, but no source of UV radiation is known within the human body.

Recently, the proposal [5] was made to use nanoparticles (NPs) to attach to the virus in combination with IR light to cause a structural change that stops the ability of the virus to survive and reproduce. However, the structural change necessary to produce UV radiation to disinfect the Coronavirus is not identified.

### 3. Purpose

The purpose of this paper is to present the simple QED theory [6] of nanoscale heat transfer to explain how NPs produce UV radiation within the human body to disinfect the Covid-19 virus. Developed over the past decade, simple QED in Covid-19 treatment is implemented by *in vivo* vaccinations of NPs in saline.

However, UV radiation also damage brain neurons and DNA, but may be justified in the brief time necessary to disinfect the Coronavirus. Biodegradable lipid NPs avoid long term UV damage. The UVC radiation emission is powered by heat of the blood at body temperature, the simplicity of which is illustrated in Fig. 1.




Figure 1. UVC disinfection of Covid-19 by nanoparticles

### 4. Background

Simple QED is a nanoscale heat transfer process based on the Planck law [7] of quantum mechanics (QM) differing significantly from that of classical physics. Research in nanoscale heat transfer [8-10] has advanced over the past decades, and a large number of interesting phenomena have been reported. But despite the research advances, there are still challenges remaining in understanding the mechanism of nanoscale thermal transport. Perhaps this is so, because researchers have not appreciated the significant difference between classical physics and the Planck law with regard to the heat capacity of the atom, without which nanoscale heat transfer cannot proceed.

In this regard, the Planck law denies atoms in nanostructures the heat capacity to change in temperature upon the absorption of heat - a difficult notion to accept because of training in classical physics. Heat transfer without changes in temperature preclude the Fourier law of heat conduction commonly used in nanoscale heat transfer. Similarly, the Stefan-Boltzmann law for radiative heat transfer depending on temperature is not applicable to nanostructures. Although valid at the macroscale, the Fourier law and Stefan-Boltzmann equation are invalid at the nanoscale. Molecular Dynamics (MD) simulations [11] based on classical physics thought to provide an understanding of the atomic response to thermal disturbances assume atoms in nanostructures have temperature. Although MD is valid for periodic boundary conditions, extension to

discrete nanostructures is not. Researchers need both new theory and computational procedures to be developed to understand nanoscale heat transfer.

Simple QED is a method of nanoscale heat transfer analysis that conserves heat with EM radiation instead of temperature. QED stands for quantum electrodynamics, a complex theory based on *virtual* photons advanced by Feynman [12] and others. In contrast, simple QED is a far simpler theory based on the Planck law that requires the heat capacity of the atoms in nanostructures to vanish allowing conservation to proceed by the creation of *real* photons comprising EM waves that stand within and across the nanostructure. Unlike electron level quantum states, simple QED quantum states are size dependent based on the dimension of the nanostructure over which the EM waves stand.

By classical physics, the  $kT$  heat capacity of the atom is independent of the EM confinement wavelength  $\lambda$ , where  $k$  is the Boltzmann constant and  $T$  absolute temperature. QM differs as the  $kT$  heat capacity of the atom decreases under EM confinement  $\lambda < 200$  microns, and at the nanoscale for  $\lambda < 100$  nm, the heat capacity may be said to vanish. The Planck law at 300 K is illustrated in Fig. 2.




Figure. 2: Planck law of the Atom at 300 °K

In the inset,  $E$  is Planck energy,  $h$  Planck's constant,  $c$  light speed,  $k$  Boltzmann's constant,  $T$  temperature, and  $\lambda$  the EM confinement wavelength

EM confinement occurs by the high surface-to-volume ( $S/V$ ) ratio of nanostructures that requires the heat  $Q$  to almost totally be confined in the surface, the surface heat itself providing the brief EM confinement necessary to create EM waves standing across the internal dimension  $d$  of the nanostructure. For heat (or light) having wavelength  $\lambda < d$ , the absorption occurs over the incident face of the particle. However, for small particles  $d \ll \lambda$ , the light fully immerses the particle to be absorbed uniformly over the full surface. Fig. 3 illustrates  $\lambda \gg d$  with light (yellow) immersing the NP (white) and absorbed in penetration depth  $\delta$  over the full NP surface.




Figure 3. Heat Q (or light) absorbed in full NP surface

Confinement of the light Q while creating the UVC standing wave requires EM confinement at least equal to the Planck energy E of the light. The pressure P acting on the surface is given for bulk modulus B and volume strain  $\Delta V/V$  by,  $P = B \cdot \Delta V/V = 6 \cdot \delta \cdot B/d$ . But  $P = E/V = 6E/\pi d^3$  giving  $\delta = Q/\pi B d^2$ . For an 80 nm lipid NP with bulk modulus  $B \sim 2 \times 10^9 \text{ N/m}^2$ , the absorption depth  $\delta$  of a single UVC photon is  $\delta \sim 20 \text{ fm}$  - a small but necessary depth to confine the absorbed heat  $Q = E$  to the geometry of the standing wave.

Simple QED absorbs heat Q in the NP surface given by the penetration  $\delta$  depth. Unable to conserve the surface heat by a change in temperature, conservation requires the creation of simple QED radiation, the time  $\tau$  to create the standing wave,  $\tau = 2d/(c/n)$ . The Planck energy  $E \sim h/\tau = hc/2nd$  depends on the refractive index n of the NP to correct for the velocity c of light within the NP. The simple QED Planck energy E is quantized by the dimension d of the NP that defines the half-wavelength  $\lambda/2$  of the nanostructure. Fig. 4 illustrates the standing EM radiation in a spherical NP of diameter d, but NP atoms still follow their quantized electron energy levels.




Figure. 4: Planck Energy of EM Radiation

In a rectangular NP with different dimensions of width, thickness, and length there are 3 simple QED quantum states corresponding to the different dimensions of the NP. However, only the minimum dimension is important as by Fermat's principle, the absorbed heat is dissipated in minimum time. Continuous variation in internal nanostructure dimensions produces a broadband spectrum of simple QED dissipated in continuous QED quantum states. Historically, the notion of size dependent quantum states is not found in the literature.

## 5. Analysis

The simple QED analysis of the thermal response of a single lipid NP in a thermal bath of tissue and blood is illustrated in Fig. 5.




Figure. 5: NP in a Thermal Bath

The NP absorbs heat  $Q$  from the thermal bath at absolute temperature  $T$  by conduction. Fig. 2 shows Fourier's heat conduction equation at 300 K is only valid in the bath for  $kT = 0.0254$  eV. The radius  $R_s$  at which bath atoms have thermal  $kT$  energy is  $\lambda > 200$  microns. 1. For body tissue and water having refractive index  $n = 1.4$ , the radius  $R_s = \lambda/4n \sim 36$   $\mu\text{m}$ . What this means is the heat flow  $Q$  from the bath at temperature  $T$  is converted at  $R_s$  to EM radiation in the far IR ( $\lambda = 200$  microns) and upon being absorbed at by the NP is conserved by emitting simple QED radiation. Small temperature changes occur for  $R < 10$  microns, but clearly vanish for NPs  $< 100$  nm.

Classically, all atoms in the NP at equilibrium have temperature  $T$  equal to the bath temperature. In terms of the Boltzmann constant  $k$  and the number  $N$  of atoms, the total LNP thermal energy  $U$  is,

$$U = \frac{3}{2} NkT$$

However, by the Planck law the  $N$  atoms do not have  $kT$  energy. Instead, simple QED conserves the energy  $U$  that otherwise would occupy the 80 nm LNP by creating standing EM radiation across the NP diameter  $d$  as shown in Fig. 4.

The molecular weight of the lipid meibomian  $\text{C}_{44}\text{H}_{56}\text{O}_2$  is 616 and the number  $N_m$  of molecules is,  $N_m = (\rho V/616) \cdot A_v$ , where volume  $V = \pi d^3/6 = 2.68 \times 10^{-22}$   $\text{m}^3$ , density  $\rho = 1000$   $\text{kg}/\text{m}^3$  and Avagadro's number  $A_v = 6.023 \times 10^{26}$  mols/kg-mol. Hence,  $N = 2.62 \times 10^5$  and  $U \sim 10.8$  keV. For  $E = hc/\lambda$  at  $\lambda = 254$  nm,  $E \sim 4.88$  eV and the lipid NP creates about 2200 UVC photons equilibrating with the 300 °K thermal bath temperature. Once created, the emitted UVC photons are absorbed by the Covid-19 virus or water bath, the bath temperature  $T$  once again produces the number of 2200 UVC photons repetitively.

But how rapidly does the NP surface temperature recover?

The simple QED creation of UVC having Planck energy  $E = 4.88$  eV absorbing a pulse of heat from the water changing the temperature  $\Delta T$  given [14] by,

$$\Delta T = \frac{1.2}{\pi d^2 K} \left( \frac{E}{\Delta t} \right) \sqrt{\frac{\alpha}{\pi}} [\sqrt{t + \Delta t} - \sqrt{t}]$$

The Planck energy  $E$  is spread over the spherical surface area  $\pi R_s^2$ . The pulse duration is  $\Delta t = 2d / (c/n) \sim 0.85$  fs. The UVC heat  $Q = E/\Delta t \sim 900$   $\mu$ W. For water, thermal diffusivity  $\alpha = K/\rho C$ , where  $\alpha = 1.24 \times 10^{-7}$   $m^2/s$  and  $K = 0.52$  W/m-K. Fig. 6 shows the initial temperature  $\Delta T$  to be an imperceptible  $\sim 2$   $\mu$ °C that recovers in  $< 1$  ps.




Figure. 6: Single Photon Creation Time

However, the UVC photon must be created promptly, say  $< 5$  fs as noted by the blue circle. What this means is the UVC photon cannot be created from body temperature surroundings. Much higher bath temperatures are required. To create the UVC photon, the required temperature  $T = E/1.5k \sim 37,000$  °K. But high temperatures are not necessary for NPs under high EM confinement. Indeed, once the heat  $Q$  is absorbed in the penetration depth  $\delta = 20$  fm of the NP, the creation time  $\tau$  of the UVC photon  $\tau = 2d/(c/n) \sim 0.85$  fs  $< 5$  fs and acceptable.

## 6. Conclusions

In simple QED, the Planck law allows the lipid NPs to produce UVC radiation that disinfects the Covid-19 from heat at body temperature, a significant difference with classical physics that predicts the lipid NP only acquires the temperature of the bath.

External lasers are not required to excite the NPs. Simple QED produces UVC from lipid NPs using only the thermal energy of the surrounding blood and tissue.

In the manner of an *in vivo* Covid-19 vaccine, the NP treatment of UVC disinfection kills the live virus to produce the inactivated virus that acts as the antigen to elicit immunity to subsequent Covid-19 infection.

The CDC is requested to develop UVC treatment of Covid-19 by *in vivo* vaccination doses of  $\sim 80$  nm lipid NPs. In the blood stream, the disinfection would be rapid. Doses should be minimal as NPs entering the brain damage neurons and DNA. Nevertheless, for patients in a life-threatening condition, the attendant brain damage would appear justified.

## References

1. Reed NG. (2010). The History of Ultraviolet Germicidal Irradiation for Air Disinfection. *Public Health Reports*, 125: 15-27.
2. Downes A, Blunt TP. (1877). Researches on the effect of light upon bacteria and other organisms. *Proc R Soc Lond.* 26:488-500.
3. Tyndall J. (1878). Note on the influence exercised by light on organic infusions. *Proc R Soc Lond.* 28:212-3.
4. Hollaender A, Oliphant JW. (1944). The inactivating effect of monochromatic ultraviolet radiation on influenza virus. *J Bacteriol.* 1948:447-54.
5. Candanos RM. (2020). Here's how nanoparticles could help us get closer to a treatment for COVID-19. See <https://phys.org/news/2020-03-nanoparticles-closer-treatment-covid-.html>
6. Prevenslik T. See Simple QED Applications. See [www.nanoqed.org](http://www.nanoqed.org) at 2010-2020.
7. Planck M.: On the Theory of the Energy Distribution Law of the Normal Spectrum. *Verhandl. Dtsch. Phys. Ges.* 1900; 2: 2-37.
8. Cahill DG, et al. Nanoscale thermal transport. *Appl. Phys.* 2003; 93:793–818.
9. Chen G. Nanoscale energy transport and conversion: parallel treatment of electrons, molecules, phonons, and photons. Oxford University Press, 2005.
10. Cahill DG, et al. Nanoscale thermal transport. II. 2003–2012. 2014; *Appl. Phys. Rev.* 1, 011305.
11. Poulidakos D, Arcidiacono S, Muruyama S. Molecular Dynamics simulation in nanoscale Heat Transfer: a review. *Microscale Thermophysical Engineering.* 2003; 7:181–206.
12. Feynman R. *QED: The Strange Theory of Light and Matter.* Princeton University Press, 1976.
13. Tiffany JM. (1986) Refractive index of meibomian and other lipids. *Current Eye Research - Short Communication.* 5:887-889.
14. Carslaw HS, Yeager JC. *Conduction of Heat in Solids*, 2 nd. Ed. Oxford Univ. Press, 1959.